

Quantum Probabilistic Subroutines and Problems in Number Theory

A. Carlini and A. Hosoya

Department of Physics, Tokyo Institute of Technology, Oh-Okayama, Meguro-ku, Tokyo 152, Japan

We present a quantum version of the classical probabilistic algorithms à la Rabin. The quantum algorithm is based on the essential use of Grover's operator for the quantum search of a database and of Shor's Fourier transform for extracting the periodicity of a function, and their combined use in the counting algorithm originally introduced by Brassard et al. One of the main features of our quantum probabilistic algorithm is its full unitarity and reversibility, which would make its use possible as part of larger and more complicated networks in quantum computers. As an example of this we describe polynomial time algorithms for studying some important problems in number theory, such as the test of the primality of an integer, the so called 'prime number theorem' and Hardy and Littlewood's conjecture about the asymptotic number of representations of an even integer as a sum of two primes.

PACS numbers: 03.67.Lx, 89.70.+c, 02.10.Lh

I. INTRODUCTION

Quantum computers allow a superposition of $|0\rangle$ and $|1\rangle$ qubits with coefficients being complex numbers α and β ,

$$|\psi\rangle = \alpha|0\rangle + \beta|1\rangle. \quad (1)$$

It is this superposition which provides us with an enormous number of parallel computations by generating a superposed state of a large number of terms, for example starting with the flat superposition $(|0\rangle + |1\rangle)^N$.

Quantum computers can do unitary transformations and also make quantum mechanical observations which induce an instantaneous state reduction to $|0\rangle$ or $|1\rangle$ with the probability $|\alpha|^2$ or $|\beta|^2$, respectively [1-3].

At present there are two main kinds of interesting quantum algorithms which can beat their classical counterparts, i.e. Shor's algorithm for factoring integers [4] and Grover's algorithm for the unstructured database search [5] which achieve, respectively, an exponential and square root speed up compared to their classical analogues. One of the most interesting algorithms where these two basic unitary blocks are exploited in conjunction is the counting algorithm introduced by Brassard et al. [6] (see section 3 for a detailed description), which can count the cardinality t of a set of states with a given property present in a flat superposition of N states in a time which is polynomial in the ratio N/t , and with an accuracy which can be made exponentially close to one.

In this work we shall show how an extended use of this algorithm can be exploited to construct unitary and fully reversible operators which are able to emulate at the quantum level a class of classical probabilistic algorithms. Classical probabilistic algorithms are characterized by the use of random numbers during the computation, and the fact that they give the correct answer with a certain probability of success, which can be usually made

exponentially close to one by repetition (see, e.g., ref. [7]). In this paper we show explicitly a series of quantum, fully reversible and unitary algorithms which can be seen as the quantum analogue of the aforementioned classical randomized algorithms, in the sense that they naturally select the 'correct' states with an arbitrarily large probability amplitude in the end of the computation, and that the final measuring process is only an option which may not be used, e.g., in the case when the 'answer' provided by such quantum algorithms is needed as a partial (subroutine) result for further computations in a larger and more complex quantum network. The main ingredients for the construction of our quantum algorithms consist in the repeated use of the quantum counting transform of ref. [6], the exploitation of the resulting interference and entanglement among quantum states and, to some extent, in assigning to some extra ancilla qubits the role analogue of the classical random repetitions. Previous work [8] (see also ref. [9] for other types of subroutines in the context of generalized quantum computation) also dealt with the problem of building unitary and reversible subroutines for the use in larger quantum computational networks. However, as quantum interference is not exploited, these methods in general require a larger memory space compared to our algorithms (see also the footnote at the end of section III).

The paper is structured as follows. In section 2 we summarize the main properties of one of the prototypes of the classical randomized algorithms, i.e. Rabin's test for primality of an integer. In section 3 we describe the main block of our quantum algorithm and, as a warm up exercise propaedeutical to section 4, we study again the case of the test of primality for a given integer, comparing our results with the classical ones. In section 4 we extend our quantum methods to the problem of checking the so called 'prime number theorem' concerning the distribution of primes smaller than a given integer. We conclude in section 5 with some discussion and future per-

spectives. Finally, in the appendix we suggest how one might test Hardy and Littlewood's formula concerning the asymptotic behaviour of the number of representations of a given even integer as the sum of two primes, and also comment about the possible proof of a famous Goldbach's conjecture in number theory.

II. CLASSICAL RANDOMIZED PRIMALITY TEST

One of the prototype examples of classical probabilistic algorithms is that of Rabin [7] for testing the primality of a given number k .^{*} The algorithm is probabilistic as it uses random integers during the computation, it is always correct when certifies a number to be composite, while it asserts primality with an arbitrarily small probability of error. The algorithm tests the following condition $W_k(a)$, for $1 \leq a < k$ (and factoring out of $k-1$ the highest power of 2 which divides it, i.e. writing $k-1 \equiv 2^h l$, with h integer and l odd):

- (i) $a^{k-1} \bmod k \neq 1$;
 - (ii) $\exists i \in [1, h] / \gcd(a^{(k-1)/2^i}, k) \neq 1$.
- (2)

If at least one of conditions (i) or (ii) is satisfied, then $W_k(a) = 0$ and a is said to be a witness to the compositeness of k . On the other hand, if neither (i) nor (ii) are satisfied, then $W_k(a) = 1$. The most important property of the witness function $W_k(a)$ is that for a composite number $k \equiv k_C$ the number t_{k_C} of witnesses a s.t. $W_{k_C}(a) = 0$ is

$$t_{k_C} \geq \frac{3(k_C - 1)}{4}, \quad (3)$$

i.e., for a composite k_C it is guaranteed that at least $3/4$ of the $a < k_C$ are witnesses to k_C [7]- [11] (see also ref. [12]). For a prime number $k \equiv k_P$, instead, none of the a is a witness (i.e. $W_{k_P}(a) = 1$ for all $1 \leq a < k_P$). Conversely, if, for an integer $1 \leq a < k$ picked at random, one finds that $W_k(a) = 1$, then one can correctly declare k to be prime with a probability $3/4$, while, if one finds $W_k(a) = 0$, then k can be declared composite with certainty.

The classical randomized algorithm for the test of primality of k heavily relies on this property of the witness function W_k and proceeds as follows. Given the number k to be tested, one first picks up randomly h numbers a_i s.t. $1 \leq a_i < k$ ($i \in [1, h]$) and checks their witness function $W_k(a_i)$. If $W_k(a_i) = 0$ for *at least one* of the a_i ,

^{*}For a review of other probabilistic and deterministic classical tests of primality see, e.g., ref. [10].

then k is declared *composite*, while if $W_k(a_i) = 1$ for *all* a_i , then k is declared *prime*. If k is declared composite the test is always correct, but if k is declared prime the test may fail with a probability (i.e., the probability of independently picking h 'false' witnesses) smaller than $1/2^{2h}$ [7]. The computational complexity S_{class} of the algorithm, defined as its running time as a function of the number of required operations, is polynomial in the number of digits of k , i.e. $S_{class} \simeq O[h \text{ poly}(\log k)]$.[†]

We stress once more the main point leading to the good performance of the classical randomized algorithm, i.e. the *large gap between the number of witnesses in the cases when k is a prime and when it is a composite*.

III. QUANTUM PRIMALITY TEST

Let us now present, at first, also as a useful 'warm-up' exercise, a quantum algorithm to test the primality of a given number k and compare it with the classical probabilistic one by Rabin. The main idea underlying our quantum computation is the repeated use of the counting algorithm COUNT originally introduced by Brassard et al. [6]. The algorithm COUNT makes an essential use of two of the main tools in quantum computation, i.e. Grover's unitary operation G for extracting some elements from a flat superposition of quantum states, and Shor's Fourier operation F for extracting the periodicity of a quantum state. Grover's unitary transformation is given by $G = -WS_0WS_1$, where the Walsh-Hadamard transform W is defined as

$$W|a\rangle \equiv \frac{1}{\sqrt{k}} \sum_{b=0}^{k-1} (-1)^{a \cdot b} |b\rangle \quad (4)$$

(with $a \cdot b \equiv \sum_i a_i b_i \bmod 2$, $a_i(b_i)$ being the binary digits of $a(b)$), $S_0 \equiv I - 2|0\rangle\langle 0|$ and $S_1 \equiv I - 2 \sum_w |w\rangle\langle w|$, which changes sign to the searched states $|w\rangle$.[‡] Shor's operation is, instead, given by the Fourier transform[§]

$$F|a\rangle \equiv \frac{1}{\sqrt{k}} \sum_{b=0}^{k-1} e^{2i\pi ab/k} |b\rangle. \quad (5)$$

[†]One has to randomly generate (e.g., tossing $O[\log k]$ coins) h numbers, taking $O[\log k]$ steps for each number, and to evaluate the witness function for each of these numbers, taking another $O[\text{poly}(\log k)]$ steps [7].

[‡]From here onwards, for simplicity, we use the compact notation according to which, e.g. assuming $b \simeq O(k)$, we have $b = 2^j b_j + 2^{j-1} b_{j-1} + \dots + 2^0 b_0$ (with $j = \lceil \log k \rceil$) and $|b\rangle$ itself is actually an acronym for the tensor product of $j+1$ qubits, i.e. $|b\rangle \equiv |b_j\rangle \otimes |b_{j-1}\rangle \otimes \dots \otimes |b_0\rangle$.

[§]Note that one can write the flat superposition as $W|0\rangle = F|0\rangle = \sum_a |a\rangle / \sqrt{k}$.

Then, the COUNT algorithm can be summarized by the following sequence of operations:

COUNT:

- 1) $(W|0\rangle)(W|0\rangle) = \sum_m |m\rangle \sum_a |a\rangle$
- 2) $\rightarrow (F \otimes I)[\sum_m |m\rangle G^m(\sum_a |a\rangle)]$
- 3) $\rightarrow \text{measure } |m\rangle$

Since the amplitude of the set of the states $|w\rangle$ after m iterations of G on $|a\rangle$ is a periodic function of m , the estimate of such a period by use of the Fourier analysis and the measurement of the ancilla qubits in $|m\rangle$ will give information on the size t of this set, on which the period itself depends. The parameter P determines both the precision of the estimate t and the computational complexity of the COUNT algorithm (which requires P iterations of G).

Our quantum algorithm for the test of primality makes essential use of the COUNT algorithm for estimating the number of witnesses to the compositeness of k , and of $R \log P$ ancilla qubits $|m_i\rangle_i$ (with $m_i \in [0, P]$, $i \in [1, R]$ and P is an integer power of 2 to be determined later) which are finally measured and which are necessary in order to sharpen the constructive interference of the 'good' states.

We start with the tensor product of R $|0\rangle_i$ states with $\log P$ qubits, and one state $|0\rangle$ with $\log k$ qubits, i.e.

$$|\psi_0\rangle \equiv |0\rangle_1 \dots |0\rangle_R |0\rangle \quad (6)$$

and then act on each of these states with a Walsh-Hadamard transform W in order to obtain

$$|\psi_1\rangle \equiv \frac{\sum_{m_1=0}^{P-1} |m_1\rangle_1}{\sqrt{P}} \dots \frac{\sum_{m_R=0}^{P-1} |m_R\rangle_R}{\sqrt{P}} \frac{\sum_{a=0}^{k-1} |a\rangle}{\sqrt{k}}. \quad (7)$$

We then act on the last $|a\rangle$ state with the $|m_1\rangle_1$ $|m_R\rangle_R$ -controlled Grover operation G^m s.t.

$$|\psi_2\rangle \equiv \frac{\sum_{m_1=0}^{P-1} |m_1\rangle_1}{\sqrt{P}} \dots \frac{\sum_{m_R=0}^{P-1} |m_R\rangle_R}{\sqrt{P}} \times \frac{\sum_{a=0}^{k-1} G^{m_1+\dots+m_R} |a\rangle}{\sqrt{k}}, \quad (8)$$

where in G we use $S_1 \equiv I - 2 \sum_{W_k(a)=0} |a\rangle\langle a|$, which changes sign to the witnesses of the compositeness of k .

**

**A unitary transformation representing the witness function $W_k(a)$ can be easily obtained by defining the quantum

In the following we will assume that P is at most $\simeq O[\text{poly}(\log k)]$, so that the steps required to compute the repeated Grover operations $^{\dagger\dagger} G^{m_1+\dots+m_R}$ is polynomial in $\log k$.

We define the quantities

$$\sin \theta_k \equiv \sqrt{\frac{t_k}{k}} \quad (9)$$

and

$$\begin{aligned} k_{m_1\dots m_R} &\equiv \sin[2(m_1 + \dots + m_R) + 1]\theta_k \\ l_{m_1\dots m_R} &\equiv \cos[2(m_1 + \dots + m_R) + 1]\theta_k, \end{aligned} \quad (10)$$

where t_k is the number of a s.t. $1 \leq a < k$ and $W_k(a) = 0$, and the states

$$\begin{aligned} |B_1\rangle_k &\equiv \frac{1}{\sqrt{t_k}} \sum_{W_k(a)=0} |a\rangle \\ |B_2\rangle_k &\equiv \frac{1}{\sqrt{k-t_k}} \sum_{W_k(a)=1} |a\rangle, \end{aligned} \quad (11)$$

such that we can simplify eq. (8) by use of

$$\begin{aligned} \frac{1}{\sqrt{k}} \sum_{a=0}^{k-1} G^{m_1+\dots+m_R} |a\rangle &= k_{m_1\dots m_R} |B_1\rangle_k \\ &+ l_{m_1\dots m_R} |B_2\rangle_k. \end{aligned} \quad (12)$$

Next we apply Shor's Fourier transform on each of the R ancilla states $|m_i\rangle_i$ in order to extract the periodicity θ_k (and, therefore, via eq. (9), the number of witnesses t_k) which is hidden in the amplitudes $k_{m_1\dots m_R}$ and $l_{m_1\dots m_R}$, i.e. we transform $|\psi_2\rangle$ into

AND of the basic operations in (i) and (ii) in eq. (2), each of which can be evaluated in a time which is polynomial in $\log k$. For instance, one can first evaluate h in $k-1 \equiv 2^h l$ by reading the highest qubit in $k-1$, and then build the state $|W_k^{(0)}(a)\rangle \dots |W_k^{(h)}(a)\rangle |W_k(a)\rangle$, where $W_k^{(i)}(a) \equiv \Theta[a^{k-1} \bmod k]$, for $i \in [1, h]$ we have $W_k^{(i)}(a) \equiv \Theta[\text{GCD}(a^{(k-1)/2^i}, k) - 1 \bmod (k-1) + 1]$, with $\Theta[1] = 1$ and $\Theta = 0$ otherwise, and where $|W_k(a)\rangle \equiv |W_k^{(0)}(a)\rangle \text{ AND } \dots W_k^{(h)}(a)\rangle$. The operator $S_1 \sum_a |a\rangle = -\sum_a (-1)^{W_k(a)} |a\rangle$ can then be easily realized by tensoring the states $|a\rangle$ with the ancilla qubit $|e\rangle \equiv [|0\rangle - |1\rangle]/\sqrt{2}$ and acting with $U_{W_k} : |a\rangle |e\rangle \rightarrow |a\rangle |e + W_k(a) \bmod 2\rangle$. All the operations leading to the evaluation of $W_k(a)$, except the last for the phase change, have to be undone again, as usual, before acting with S_1 and G .

†† Each of which has a computational complexity $S_G \simeq O[\text{poly}(\log k) + \log k] \simeq O[\text{poly}(\log k)]$, the first term being for the quantum parallel evaluation of the witness functions, and the second term for the evaluations of the W and S_0 transforms.

$$\begin{aligned}
|\psi_3\rangle &\equiv \frac{\sum_{m_1, l_1=0}^{P-1} e^{2i\pi l_1 m_1/P} |l_1\rangle_1 \dots}{P} \\
&\times \frac{\sum_{m_R, l_R=0}^{P-1} e^{2i\pi l_R m_R/P} |l_R\rangle_R}{P} \\
&\times [k_{m_1 \dots m_R} |B_1\rangle_k + l_{m_1 \dots m_R} |B_2\rangle_k]. \quad (13)
\end{aligned}$$

After some elementary algebra, eq. (13) can be rewritten as

$$\begin{aligned}
|\psi_3\rangle &\equiv \frac{1}{2} \sum_{l_1, \dots, l_R=0}^{P-1} |l_1\rangle_1 \dots |l_R\rangle_R e^{-i\pi(l_1 + \dots + l_R)P} \\
&\times \left[e^{i\pi f_k^{(R)}} \prod_{i=1}^R s_{l_i+}^{(P)} (-i|B_1\rangle_k + |B_2\rangle_k) \right. \\
&\left. + e^{-i\pi f_k^{(R)}} \prod_{i=1}^R s_{l_i-}^{(P)} (i|B_1\rangle_k + |B_2\rangle_k) \right], \quad (14)
\end{aligned}$$

where we have introduced the following quantities,

$$\begin{aligned}
f_k &\equiv \frac{P\theta_k}{\pi} \quad ; \quad 0 \leq f_k \leq \frac{P}{2} \\
f_k^{(R)} &\equiv f_k \left[R + \frac{(1-R)}{P} \right] \quad (15)
\end{aligned}$$

and

$$s_{l_i \pm}^{(P)} \equiv \frac{\sin \pi(l_i \pm f_k)}{P \sin \frac{\pi(l_i \pm f_k)}{P}}. \quad (16)$$

In particular, when counting the witnesses for a given k , we have two different possibilities: either k is a prime, in which case we have that $t_{k_P} = 0$ and therefore $\theta_{k_P} = f_{k_P} = 0$; or k is a composite, for which we have that $t_{k_C} \geq 3k_C/4$ and $\theta_{k_C} \geq \pi/3$, implying that $P/3 \leq f_{k_C} \leq P/2$.

Going back to eq. (14), we can see that, in the case when k is a prime, G effectively acts as an identity operator, so that $|\psi_3\rangle$ simplifies to

$$|\psi_3\rangle \rightarrow |0\rangle_1 \dots |0\rangle_R |B_2\rangle_k \quad ; \quad \text{when } k = k_P. \quad (17)$$

On the other hand, when k is a composite, almost all of the ancilla qubits in $|\psi_3\rangle$ will be in a state different from $|0\rangle_1 \dots |0\rangle_R$. In fact, the probability of finally measuring $|0\rangle_1 \dots |0\rangle_R$ when k is composite is

$$\begin{aligned}
P(|0\rangle_1 \dots |0\rangle_R) &\Big|_{k_C} = (\alpha_k)^{2R} \Big|_{k_C} \equiv \left(\frac{\sin \pi f_k}{P \sin \frac{\pi f_k}{P}} \right)^{2R} \Big|_{k_C} \\
&\leq \left(\frac{2}{\sqrt{3}P} \right)^{2R} \simeq O[P^{-2R}], \quad (18)
\end{aligned}$$

since we have $f_{k_C} \geq P/3$.

Summarizing the above results, our quantum algorithm for testing the primality of a given number k is

probabilistic in the following sense: if in the final measurement process of the $R \log P$ ancilla qubits we obtain a state with *at least one* of the qubits different from $|0\rangle$, we can declare with *certainly* that the number k is a *composite*; on the other hand, if *all* the ancilla qubits are in the state $|0\rangle$, we can claim with an *error probability smaller than* $O[P^{-2R}]$ that the number k is a *prime*.

The use of $R \log P$ ancilla qubits and the repeated application of the Fourier transforms is made in order to sharpen the constructive interference effects at the basis of the measurement of the period θ_k and, in this sense, it can be seen as the quantum analogue of the multiple random tests used in the classical primality algorithm by Rabin. Moreover, our algorithm, provided that the final measurement step is omitted, is clearly unitary and fully reversible, and as such it can be used as an intermediate unitary transform inside a larger and more complicated algorithm. The strength of this quantum algorithm then critically relies on the use of the superposition and entanglement of states, and above all on the existence of a *gap* between the cardinalities of different sets of the domain of a given test function (in the case of the function $W_k(a)$ the domain is divided in the set of states with $W_k = 1$, $f_k = 0$, and those with $W_k = 0$, $f_k \geq P/3$).^{‡‡} We will show other and more interesting problems where these properties can be fruitfully exploited in section 4 and in the appendix.

The computational complexity of the quantum algorithm can be written as $S_{quant} \simeq O[\log k + R\{\log P + (\log P)^2 + PS_G\}] \simeq O[RPS_G]$,^{§§} with the number of steps required for G given by $S_G \simeq O[\text{poly}(\log k)]$ (see footnote below eq. (8)), so that we obtain $S_{quant} \simeq O[R \text{poly}(\log k)]$.^{***}

^{‡‡}Taking $f_{good} = 0$ (i.e., $\alpha_{good} = 1$) and, more in general, $f_{bad} = \xi P/2$ with $\xi \in (0, 1]$, and requiring a success probability exponentially close to one, we have to choose, e.g., $\alpha_{bad} \leq (P \sin \pi \xi/2)^{-1} < 1/2$ in eq. (18). Thus, actually even small values of ξ (but $\xi > [\text{poly}(\log N)]^{-1}$) are good enough for the quantum test to be sufficiently reliable, provided one takes $P \geq O[\xi^{-1}]$.

^{§§}The first term is for the construction of the flat superposition $\sum_a |a\rangle$, while the other terms arise, in the order from left to right, from the evaluations of the R flat superpositions of the ancilla states $\sum_{m_i} |m_i\rangle_i$ ($i \in [1, R]$), the R Fourier transforms on the same states and the operation $G \sum^{m_i}$, requiring PR repetitions of the basic block G .

^{***}One might observe that, in fact, it is not necessary to use COUNT for primality testing, but simply build the state $[\sum_a |a\rangle |W_k(a)\rangle]$ (for the bases a of the integer k), measure the ancilla qubit $|W_k(a)\rangle$ and repeat the procedure h times. This would imply the same error probability $P_e \simeq 2^{-2h}$ (after h trials), computational complexity $S_{quantum} \simeq O[\text{poly}(\log k)]$ and memory space required $M = \log k + 1$ as

IV. COUNTING $K_P < N$ AND THE PRIME NUMBER THEOREM

One of the problems in which the quantum algorithm of the previous section can be explicitly used, as the basic block of another more complex unitary operation, is the case of the testing of the so called 'prime number theorem' (see, e.g., ref. [10] and references therein), according to which the total number t_N of primes k_P smaller than a given number N is given by the formula

$$t_N \equiv \pi(N) \simeq \frac{N}{\log N} \quad (19)$$

Our quantum algorithm essentially consists of a sub-loop which checks for the primality of a given $k < N$ by counting its witnesses, a main loop for the counting of primes less than N , and a final measurement of some ancilla qubits. More in details, we can schematically summarize the main operations in the following steps:

MAIN-LOOP:

Count $\#\{k | k = k_P < N\}$ using **COUNT** with $G \rightarrow \tilde{G}$ and $S_1 \rightarrow \tilde{S}_1 \equiv 1 - 2 \sum_{k_P} |k_P\rangle\langle k_P|$ (parameter Q)

SUB-LOOP:

Parallel primality tests $\forall k < N$ (parameter P) and (approximate) construction of $\tilde{S}_1$

Let us start from the SUB-LOOP of the algorithm first. The unitary transform $\tilde{S}_1$ to be computed should approximate with a high level of accuracy the following basic operation

$$\tilde{S}_1 : \frac{1}{\sqrt{N}} \sum_{k=0}^{N-1} |k\rangle \rightarrow \frac{1}{\sqrt{N}} \sum_{k=0}^{N-1} (-1)^{F_k} |k\rangle, \quad (20)$$

in our algorithm with the choice $R = 1$, $P \simeq O(1)$ repeated h times, and in the classical algorithm of ref. [7]. However, this method of repeated trials would not make the subroutine for primality testing unitary and reversible. To achieve this goal, one might instead consider (see, e.g., ref. [8]) the initial state $[\sum_{a_1} |a_1\rangle |W_k(a_1)\rangle] \dots [\sum_{a_R} |a_R\rangle |W_k(a_R)\rangle]$ and parallelly check for the R ancilla qubits $|W_k(a_i)\rangle$, for which the error probability would be $P_{e,bennett} \simeq 2^{-2R}$, the computational complexity would be $S_{quantum,bennett} \simeq O[R(poly(\log k))]$ and the memory space used would be $M_{bennett} = R(\log k + 1)$. With the choice $P \simeq O(1)$ and $R \simeq O(\log k)$, however, our algorithm has the same error probability and computational complexity, but requires logarithmically less memory space. This makes explicit the advantage of exploiting the interference among quantum states which is inherent in our method via the use of COUNT.

where $F_k \equiv 1$ for a prime $k = k_P$ and $F_k \equiv 0$ for a composite $k = k_C$.

In order to construct such an $\tilde{S}_1$, we start from the flat superposition of states $|k\rangle$ tensored with two ancilla states $|0\rangle_P$ and $|0\rangle_k$ with, respectively, $\log P$ and $\log k$ qubits, i.e.

$$|\bar{\psi}_0\rangle \equiv \frac{1}{\sqrt{N}} \sum_{k=0}^{N-1} |k\rangle |0\rangle_P |0\rangle_k, \quad (21)$$

and we act on the first two states ($|k\rangle$ and $|0\rangle_P$) in eq. (21) with an F transform and a $|k\rangle$ -controlled F operation, respectively, to get

$$|\bar{\psi}_1\rangle \equiv \frac{1}{\sqrt{N}} \sum_{k=0}^{N-1} |k\rangle \frac{\sum_{m=0}^{P-1} |m\rangle_P \sum_{a=0}^{k-1} |a\rangle_k}{\sqrt{P} \sqrt{k}} \quad (22)$$

(with $P \simeq O[poly(\log N)]$). Then, as usual, we operate with a $|m\rangle_P$ -controlled Grover transform G^m on the last ancilla states $|a\rangle_k$ followed by a Fourier transform F on $|m\rangle_P$, obtaining

$$|\bar{\psi}_2\rangle \equiv \frac{1}{\sqrt{N}} \sum_{k=0}^{N-1} |k\rangle \left[\alpha_k |0\rangle_P |A_0\rangle_k + \sum_{m=1}^{P-1} \phi_m |m\rangle_P |A_m\rangle_k \right], \quad (23)$$

where

$$\begin{aligned} |A_0\rangle_k &\equiv \sin \pi f_k |B_1\rangle_k + \cos \pi f_k |B_2\rangle_k \\ |A_m\rangle_k &\equiv [e^{-i\pi f_k} s_{m-}^{(P)} (i|B_1\rangle_k + |B_2\rangle_k) \\ &\quad + e^{i\pi f_k} s_{m+}^{(P)} (-i|B_1\rangle_k + |B_2\rangle_k)]/2 \end{aligned} \quad (24)$$

while $\sin \theta_k$, $|B_{1,2}\rangle_k$, f_k , $s_{m\pm}^{(P)}$ and α_k have been defined, respectively, in eqs. (9), (11), (15), (16) and (18), and the phase $\phi_m \equiv \exp[i\pi m(1 - 1/P)]$.

We now act with the phase change operator S_0 on the first ancilla state $| \cdot \rangle_P$, and then undo again all the previous operations (F , G^m , and the two initial F s) finally obtaining the state

$$|\bar{\psi}_3\rangle \equiv \frac{1}{\sqrt{N}} \sum_{k=0}^{N-1} |k\rangle [|0\rangle_P |0\rangle_k - 2|C_1\rangle_{P,k}], \quad (25)$$

with

$$\begin{aligned} |C_1\rangle_{P,k} &\equiv \alpha_k \sum_{n,r=0}^{P-1} \sum_{b=0}^{k-1} e^{-2i\pi nr/P} [(Im Z_{k,r}) B_{1,k}(b) \\ &\quad + (Re Z_{k,r}) B_{2,k}(b)] \frac{|n\rangle_P |b\rangle_k}{P\sqrt{k}} \end{aligned} \quad (26)$$

and

$$\begin{aligned}
B_{1,k}(b) &\equiv \frac{\sum_{W_k(a)=0} e^{-2i\pi ab/k}}{\sqrt{t_k}} \\
B_{2,k}(b) &\equiv \frac{\sum_{W_k(a)=1} e^{-2i\pi ab/k}}{\sqrt{k-t_k}} \\
Z_{k,r} &\equiv e^{i\pi f_k} e^{-2i\pi f_k r/P}.
\end{aligned} \tag{27}$$

Noting the properties that $_{P,k} < C_1 |C_1 >_{P,k} = \alpha_k^2$ and, for a prime k_P , $\alpha_{k_P} = 1$, with $|C_1 >_{P,k_P} = |0 >_P$ $|0 >_{k_P}$, we can also rewrite eq. (25) as

$$\tilde{S}_1 |\bar{\psi}_0 > = |\bar{\psi}_3 > \equiv |\Psi > + |E >, \tag{28}$$

where

$$\begin{aligned}
|\Psi > &\equiv \frac{1}{\sqrt{N}} \sum_{k=0}^{N-1} (-1)^{F_k} |k > |0 >_P |0 >_k \\
|E > &\equiv -\frac{2}{\sqrt{N}} \sum_{k=k_C} |k > |C_1 >_{P,k},
\end{aligned} \tag{29}$$

which realize, as wanted, the operation $\tilde{S}_1$ of eq. (20), with the norm of the correction term $|E >$ upper bounded by

$$\begin{aligned}
\langle E | E > &= \frac{4}{N} \sum_{k=k_C} \alpha_k^2 \\
&\leq 4 \left(\frac{2}{\sqrt{3P}} \right)^2 \simeq O[P^{-2}].
\end{aligned} \tag{30}$$

Defining, in a symbolic notation, the sequence of operations

$$U_1 \equiv F [CTRL_{|m>_P}(G)] [CTRL_{|k>}(F)] F, \tag{31}$$

we have, in fact,

$$\tilde{S}_1 \equiv U_1^\dagger S_0 U_1. \tag{32}$$

Let us now consider the MAIN-LOOP of the algorithm, i.e. that counting the total number of $k_P < N$. Grover's transform $\tilde{G}$ entering this part of the algorithm can then be written as

$$\tilde{G} \equiv U_2 \tilde{S}_1 \quad ; \quad U_2 \equiv -W^{(k)} S_0^{(k)} W^{(k)}, \tag{33}$$

and with the caveat that now the operations $W^{(k)}$ and $S_0^{(k)}$ appearing in the operator U_2 of eq. (33) are acting on the states $|k >$, and that the states to be counted finally are those with $k = k_P$.

Defining as usual

$$\sin \theta_N \equiv \sqrt{\frac{t_N}{N}} \tag{34}$$

and the 'good' and 'bad' states, respectively, as

$$\begin{aligned}
|G > &\equiv \frac{\sum_{k_P} |k > |0 >_P |0 >_k}{\sqrt{t_N}} \\
|B > &\equiv \frac{\sum_{k_C} |k > |0 >_P |0 >_k}{\sqrt{N-t_N}},
\end{aligned} \tag{35}$$

we have then

$$\begin{aligned}
\tilde{S}_1 |G > &= -|G > \\
\tilde{S}_1 |B > &= |B > + \sec \theta_N |E >.
\end{aligned} \tag{36}$$

Consequently, we can derive a formula for the iteration of the operator $\tilde{G}$ acting on the state $|\bar{\psi}_0 >$, i.e.

$$\tilde{G}^m |\bar{\psi}_0 > = G^m |\bar{\psi}_0 > + |E_n >, \tag{37}$$

with

$$|E_n > \equiv \sec \theta_N \left[\sum_{j=1}^n l_{n-j} \tilde{G}^{j-1} \right] U_2 |E >, \tag{38}$$

where we have again used the variables $k_m \equiv \sin(2m+1)\theta_N$ and $l_m \equiv \cos(2m+1)\theta_N$, the formulas

$$\begin{aligned}
|\bar{\psi}_0 > &= \sin \theta_N |G > + \cos \theta_N |B > \\
|\Psi > &= -\sin \theta_N |G > + \cos \theta_N |B >
\end{aligned} \tag{39}$$

and eq. (36).

We have now all the building blocks necessary to proceed with the construction of the quantum algorithm counting the number of k_P s.t. $k_P < N$. We start from $|\bar{\psi}_0 >$ given by formula (21) and tensor it with a flat superposition of ancilla states $|m >_Q$ with $\log Q$ qubits, i.e.

$$|\bar{\psi}_4 > \equiv \frac{1}{\sqrt{Q}} \sum_{m=0}^{Q-1} |m >_Q |\bar{\psi}_0 > \tag{40}$$

(with Q an integer power of 2 to be chosen later of $O[\text{poly}(\log N)]$), then we act on $|\bar{\psi}_0 >$ with the $|m >_Q$ -controlled $\tilde{G}^m$ and with F on $|m >_Q$, getting

$$\begin{aligned}
|\bar{\psi}_5 > &\equiv \frac{1}{Q} \sum_{m,n=0}^{Q-1} e^{2i\pi mn/Q} |n >_Q [k_m |G > + l_m |B > \\
&+ |E_m >] = \frac{1}{2} \sum_{n=0}^{Q-1} e^{i\pi n(1-1/Q)} |n >_Q [e^{-i\pi f_Q} s_{n-}^{(Q)} \\
&\times (i|G > + |B >) + e^{i\pi f_Q} s_{n+}^{(Q)} (-i|G > + |B >)] \\
&+ \frac{1}{Q} \sum_{m,n=0}^{Q-1} e^{2i\pi mn/Q} |n >_Q |E_m >,
\end{aligned} \tag{41}$$

where

$$f_Q \equiv Q\theta_N/\pi \tag{42}$$

and $s_{n\pm}^{(Q)}$ are defined in eq. (16).

Now, the last step of the algorithm consists in measuring the value of the state $|\cdot\rangle_Q$ in $|\psi_5\rangle$. Using the expected estimate that $\theta_N \simeq O[1/\sqrt{\log N}]$, which gives $f_Q \simeq O[Q/\sqrt{\log N}]$, and by choosing

$$Q \simeq O[(\log N)^\beta] \quad ; \quad \beta > 1/2, \quad (43)$$

we get the ansatz $1 < f_Q < Q/2 - 1$ of ref. [6]. Then it can be easily shown, exactly as in ref. [6], that the probability W to obtain any of the states $|f_- \rangle_Q$, $|f_+ \rangle_Q$, $|Q - f_- \rangle_Q$ or $|Q - f_+ \rangle_Q$ (where $f_- \equiv [f_Q] + \delta f$ and $f_+ \equiv f_- + 1$, with $0 < \delta f < 1$) in the final measurement is given by

$$W \geq \frac{8}{\pi^2} - |W_{E_n}|, \quad (44)$$

where W_{E_n} is the contribution coming from terms involving $|E_n\rangle$, and whose explicit form we omit here for simplicity.

Using the upper bound $\langle E_n | E_n \rangle \leq O[n^2] \langle E | E \rangle$ and choosing

$$P \simeq O[(\log N)^\gamma] \quad ; \quad \gamma > \beta, \quad (45)$$

from eq. (30) we then get the estimate^{††}

$$\langle E_n | E_n \rangle \leq O[(\log N)^{-2(\gamma-\beta)}] \ll 1 \quad (46)$$

which, substituted in the formula for W_{E_n} and, then, in eq. (44), finally gives the estimate

$$W \geq \frac{8}{\pi^2} \{1 - O[(\log N)^{-(\gamma-\beta)}]\}. \quad (47)$$

This means that with a high probability we will always be able to find one of the states $|f_\pm \rangle_Q$ or $|P - f_\pm \rangle_Q$ and, therefore, to evaluate the number t_N from eqs. (34) and (42).^{‡‡}

Of course, as explained in ref. [6], since in general f_Q is not an integer, the measured $\tilde{f}_Q$ will not match exactly the true value of f_Q , and we will have some errors. In particular, defining $\tilde{t}_N \equiv N \sin^2 \tilde{\theta}_N$, with $\tilde{\theta}_N = \tilde{\theta}_N(\tilde{f}_Q)$, we have for the error over t_N the estimate [6]

$$\begin{aligned} |\Delta t_N|_{exp} &\equiv |\tilde{t}_N - t_N| \leq \pi \frac{N}{Q} \left[\frac{\pi}{Q} + 2\sqrt{\frac{t_N}{N}} \right] \\ &\simeq O[N(\log N)^{-\beta-1/2}]. \end{aligned} \quad (48)$$

^{†††}The condition (45) is sufficient but not necessary in order to have $|W_{E_n}| \ll 1$. In fact, one can also choose $P = cQ$, provided that the constant $c \ll 1$.

^{‡‡‡}We note that, as an alternative of choosing P as in eq. (45), one could also repeat the counting algorithm a sufficient number of times, as we did in the previous section (see eq. (7)), in order to reduce the 'error' probability W_{E_n} .

On the other hand, if we want to check the theoretical formula $t_N \equiv \pi(N)$ up to some power $\delta > 0$ in $\log N$, i.e. with

$$|\Delta t_N|_{th} \simeq O[N(\log N)^{-\delta-1}], \quad (49)$$

we have to impose that the measuring error over t_N is smaller than the precision required for testing t_N , i.e. we should have $|\Delta t_N|_{exp} < |\Delta t_N|_{th}$, which can be satisfied provided that

$$\beta > \delta + 1/2. \quad (50)$$

The computational complexity of the quantum algorithm can be written as $S_Q \simeq O[\log N + \log Q + (\log Q)^2 + Q(\log N + S_1)]$, where for the SUB-LOOP we have $S_1 \simeq O[\log N + \log P + PS_G + (\log P)^2]$, and which, using eqs. (43), (45) and $S_G \simeq O[\text{poly}(\log N)]$, finally gives the *polynomial* complexity $S_Q \simeq O[QPS_G] \simeq O[\text{poly}(\log N)]$.

As noted in ref. [6], moreover, one can further minimize the errors by successive repetitions of the whole algorithm. In particular, it is easily seen that the success probability W can be boosted exponentially close to one and an exponential accuracy can be achieved by repeating the whole algorithm many times and using the majority rule, still leaving the whole algorithm for the test of the 'prime number theorem' polynomial in $\log N$.^{§§§}

We conclude by stressing, once again, that the power of our quantum probabilistic methods essentially relies on: the *gap* between the cardinalities of the domains of the test function W_k ; the fact that the probability to obtain any of the states $f^\pm$ or $Q - f^\pm$ is bigger than 1/2 (which is true provided that the 'error' terms $|E_n\rangle$ have sufficiently small amplitude); that, finally, the error over the estimate of t_N is smaller than the precision we need (the last two conditions being strongly dependent on the choice of $Q, Q/P, \sqrt{k/t_k}$ and $\sqrt{N/t_N}$).

V. DISCUSSION

In this paper we have shown a method to build a quantum version of the classical probabilistic algorithms à la Rabin. Our quantum algorithms make essential use of some of the basic blocks of quantum networks known so far, i.e. Grover's operator for the quantum search of a database, Shor's Fourier transform for extracting the periodicity of a function and their combination in the counting algorithm of ref. [6]. The most important feature of our quantum probabilistic algorithms is that the

^{§§§}The average computational complexity can be further, slightly reduced by use of parallelism and anticipate measurements [6, 12].

coin tossing used in the correspondent classical probabilistic ones is replaced here by a unitary and reversible operation, so that the quantum algorithm can even be used as a subroutine in larger and more complicated networks. In particular, we described polynomial time algorithms for studying some problems in number theory, e.g. a primality test, the 'prime number theorem' and a conjecture concerning a certain distribution of couples of primes. Our quantum algorithm may also be useful for other similar tests and counting problems if there exists a classical probabilistic algorithm which somehow can guarantee a good success probability (e.g., problems related to the distribution of primes and pseudoprimes in number theory etc.). * It is well known that in a classical computation one can count, by using Monte-Carlo methods, the cardinality of a set which satisfies some conditions, provided that the distribution of the elements of such a set is assumed to be known (e.g., homogeneous). One further crucial strength and novelty of our algorithm is also in the ability of efficiently and successfully solve problems where such a knowledge or regularities may not be present.

APPENDIX A: TESTING HARDY AND LITTLEWOOD'S CONJECTURE

A very similar procedure can be followed for testing Hardy and Littlewood's conjecture [15] that the number, which we call $r_2(2N)$, of the possible representations of an even number $2N$ as the sum of two primes $k_P, l_P < 2N$ such that $k_P + l_P = 2N$, should be given by the asymptotic formula (modulo $O[\log \log N]$ factor corrections [15])

$$r_2(2N) \Big|_{th} \simeq O \left[\frac{N}{(\log N)^\mu} \right] \quad ; \quad \mu \simeq 2. \quad (A1)$$

The quantum algorithm for counting such couples and testing the conjecture can be built starting from the state

$$|\phi_0\rangle \equiv \frac{1}{\sqrt{2N}} \sum_{k=0}^{2N-1} |k\rangle |2N-k\rangle \times |0\rangle_1 |0\rangle_2 |0\rangle_3 |0\rangle_4 \quad (A2)$$

where the four ancilla states $|0\rangle_i$ have, respectively, $\log P$ qubits for $i = 1, 2$ and $\log(2N-1)$ qubits for

*Other quantum algorithms dealing with problems in number theory, such as integer factoring, finding discrete logarithms or a Pocklington-Lehmer primality test can be found, respectively, in refs. [4] and [13]. The extent to which the algorithm presented in the latter work, however, can be actually used as an efficient primality test is very questionable for us.

$i = 3, 4$, and then, as done in section 4, constructing an operator S'_1 such that

$$S'_1 : |\phi_0\rangle \rightarrow |\Phi_0\rangle + |E'\rangle, \quad (A3)$$

with the main contribution $|\Phi_0\rangle$ given by the wanted phase change for the 'good' states $|k_P\rangle > |(2N-k)_P\rangle$, i.e.

$$|\Phi_0\rangle \equiv \frac{1}{\sqrt{2N}} \sum_{k=0}^{2N-1} (-1)^{G_k} |k\rangle |2N-k\rangle \times |0\rangle_1 |0\rangle_2 |0\rangle_3 |0\rangle_4, \quad (A4)$$

where $G_k \equiv 1$ for the 'good' couples $|k_P\rangle > |(2N-k)_P\rangle$ and $G_k \equiv 0$ for all the other couples, and $|E'\rangle$ is a correction whose amplitude should be negligible with respect to that of $|\Phi_0\rangle$. This result is achieved, once again, starting from $|\phi_0\rangle$ and acting with an F on the first two ancilla states, with a $|k\rangle$ - and $|2N-k\rangle$ - 'controlled' operator F , respectively, on each of the last two ancilla states, then operating with a controlled- G transform on each of the last two ancilla states and with an F transform on each of the first two ancilla states, inverting the phase of the state $|0\rangle_1 |0\rangle_2 |0\rangle_3 |0\rangle_4$ and finally undoing the previous operations again. Doing so, one obtains as promised eqs. (A3-A4), with the explicit formula for the state $|E'\rangle$ given by (we omit all the algebraic details of the derivation for the sake of simplicity)

$$|E'\rangle \equiv -\frac{2}{\sqrt{2N}} \sum_{k'=0}^{2N-1} \alpha_{k'} \alpha_{(2N-k)'} |k'\rangle |(2N-k)'\rangle \times |C_2\rangle_{k'}, \quad (A5)$$

where the sum k' is over all couples except $|k_P\rangle > |(2N-k)_P\rangle$, and the norm of the state $|C_2\rangle_{k'}$ (a certain tensor product of the four ancilla states) is of $O(1)$.

To count the number of 'good' couples one has to repeat exactly the same steps as described from eq. (33) (with $\tilde{S}_1 \rightarrow S'_1$) to eq. (48) (with $t_N \rightarrow r_2(2N)$) in section 4, and the final result is that, again, the expected theoretical behaviour of $r_2(2N)$ can be tested up to exponential accuracy in a polynomial number of steps. In particular, if one takes

$$Q \simeq O[(\log N)^\rho] \quad ; \quad \rho > \mu/2 \\ P \simeq O[(\log N)^\sigma] \quad ; \quad \sigma > \rho, \quad (A6)$$

so that $1 < f_Q < Q/2 - 1$ in the MAIN-LOOP of the algorithm, that $\langle E'_n | E'_n \rangle \leq O[P^{-2(\sigma-\rho)}]$ and $W \geq 8/\pi^2 \{1 - O[P^{-(\sigma-\rho)}]\}$, if we want to test the $2N$ -dependence of $r_2(2N)$ with a theoretical error

$$|\Delta r_2(2N)|_{th} \simeq O[N(\log N)^{-\mu-\nu}] \quad ; \quad \nu > 0, \quad (A7)$$

the condition $|\Delta r_2(2N)|_{exp} < |\Delta r_2(2N)|_{th}$ requires that

$$\rho > \mu/2 + \nu. \quad (\text{A8})$$

The evaluation of the quantum computational complexity of the algorithm can be shown to be polynomial in the number $2N$ in a fashion similar to section 4.

Finally, we should comment that the method used could be easily extended to other similar counting problems, such as the case, e.g., when one wants to check that a given integer is the sum of more than two primes or their certain powers etc... In fact, provided that one is able to check with this quantum algorithm that *any* odd integer less than $N_0 = 10^{43000}$ (!) [16] can be written as the sum of three primes, then one might also have a numerical (although probabilistic) tool to prove with a *polynomial* computational complexity the weaker version of a famous Goldbach's conjecture, i.e. that every odd integer $N > 5$ is the sum of three primes (see, e.g., ref. [10]).

Acknowledgements

A.H.'s research was partially supported by the Ministry of Education, Science, Sports and Culture of Japan, under grant n. 09640341. A.C.'s research was supported by the EU under the Science and Technology Fellowship Programme in Japan, grant n. ERBIC17CT970007; he also thanks the cosmology group at Tokyo Institute of Technology for the kind hospitality during this work. Both authors would like to thank Prof. T. Nishino for introducing them to Rabin's primality test, Prof. N. Kurokawa for helpful discussions and the referees for pointing out some useful comments.

- [7] M.O. Rabin, Journ. Num. Th., **12**, 128 (1980); also in Algorithms and Complexity, Recent Results and New Directions, ed. J.F. Traub (Academic Press, New York, 1976), p. 21; C.L. Miller, J. Comp. Syst. Sci., **13**, 300 (1976).
- [8] C. Bennett, E. Bernstein, G. Brassard and U. Vazirani, SIAM J. Comp., **26**, 1510, (1997).
- [9] D. Aharonov, A. Kitaev and N. Nisan, Proc. 30th Annual Symposium on Theory of Computation (STOC), 20 (1997).
- [10] P. Ribenboim, The New Book of Prime Number Records (Springer-Verlag, New York, 1996).
- [11] L. Monier, Theor. Comp. Sci., **12**, 97 (1980).
- [12] I. Damgard, P. Landrock and C. Pomerance, Math. Comp., **61**, 177 (1993).
- [13] H.F. Chau and H.K. Lo, Los Alamos e-print quant-ph/9508005.
- [14] M. Luby and W. Ertel, Tech. Rep., Int. Comp. Sci. Inst., Berkeley, CA, (1993); R. Gingrich, C. Williams and N. Cerf, Los Alamos e-print quant-ph/9904049.
- [15] G.H. Hardy and J.E. Littlewood, Acta Math., **44**, 1 (1923); I.M. Vinogradov, Dokl. Akad. Nauk SSSR, **15**, 169 (1937).
- [16] J.R. Chen and Y. Wang, Acta Math. Sci. Sinica, **32**, 702 (1989).

-
- [1] P. Benioff, Journ. Stat. Phys., **22**, 563 (1980).
 - [2] D. Deutsch, Proc. Roy. Soc. London, Ser. A **400**, 96 (1985).
 - [3] R.P. Feynman, Found. Phys., **16**, 507 (1986).
 - [4] P.W. Shor, in Proceedings of the 35th Annual Symposium on Foundations of Computer Science, ed. S. Goldwater (IEEE Computer Society Press, New York, 1994), p. 124; SIAM Journ. Comput., **26**, 1484 (1997).
 - [5] L.K. Grover, in Proceedings of the 28th Annual Symposium on the Theory of Computing (ACM Press, New York, 1996), p. 212; Phys. Rev. Lett., **79**, 325 (1997).
 - [6] G. Brassard, P. Hoyer and A. Tapp, Los Alamos e-print quant-ph/9805082; M. Boyer, G. Brassard, P. Hoyer and A. Tapp, in Proceedings of the 4th Workshop on Physics and Computation, ed. T. Toffoli et al. (New England Complex Systems Institute, Boston, 1996), p. 36; also in Los Alamos e-print quant-ph/9605034 (1996) and Fortsch. Phys., **46**, 493 (1998).